

## Risks of unfavourable clinical outcomes in patients with first diagnosed stroke-associated atrial fibrillation

Zolotovskaya I., Duplyakov D., Svistunov A., Davydkin I., Osadchuk A., Osadchuk M., Trushin M.  
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

---

### Abstract

© 2018, Pharmainfo Publications. All rights reserved. Background—Atrial fibrillation (AF) is the most common arrhythmia often associated with cardioembolic stroke. Hypothesis—We investigated cases of AF first diagnosed (fdAF) in the acute phase of stroke. This type of AF might be common, and might suggest poor prognosis for patients. Methods— Following screening (n=1291) 661 patients with AF and a history of stroke were enrolled in this open-label, prospective study; patient average age was 68.0 years. Results—A high level of co-morbidity was observed: hypertension (in 100% of patients), ischemic heart disease (in 61.4% of patients), diabetes mellitus (in 52.3% of patients), and chronic kidney disease (in 32.1% of patients). In 154 (23.3%) patients AF was first diagnosed at the onset of cardioembolic stroke. The HAS-BLED score, age, hypertension and ischemic heart disease rates among these patients were comparable with the rest of the group; however, the percentage of patients with chronic kidney disease, diabetes mellitus or myocardial infarction was significantly higher. The highest one-year death rate was seen in patients with fdAF (death reported in 46 [29.9%] patients). The age of patients with fdAF who died was significantly higher, and their neurological disorders were more severe. Conclusions—The results show a high rate of fdAF and its poor prognosis in patients with cardioembolic stroke. Thus, the term “first diagnosed stroke-associated atrial fibrillation” seems relevant. The described condition has a negative predictive value, and is also associated with high co-morbidity burden in patients with fdAF.

---

### Keywords

Anticoagulant treatment, Arterial stiffness, Atrial fibrillation, Cardioembolic stroke, Endothelial dysfunction, Novel oral anticoagulants (NOACs)

### References

- [1] Boriani G, Glotzer TV, Santini M, West TM, De Melis M, Sepsi M, Gasparini M, Lewalter T, Camm JA, Singer DE (2014) Device-detected atrial fibrillation and risk for stroke: an analysis of >10,000 patients from the SOS AF project (Stroke prevention Strategies based on Atrial Fibrillation information from implanted devices). *Eur Heart J* 35(8):508-516.
- [2] Benjamin EJ, Wolf PA, D'Agostino RB, Silbershatz H, Kannel WB, Levy D (1998) Impact of atrial fibrillation on the risk of death: the Framingham Heart Study. *Circulation* 98:946-952.
- [3] Desai Y, El-Chami MF, Leon AR, Merchant FM (2017) Management of Atrial Fibrillation in Elderly Adults. *J Am Geriatr Soc* 65(1):185-193.
- [4] Chugh SS, Roth GA, Gillum RF, Mensah GA (2014) Global burden of atrial fibrillation in developed and developing nations. *Global Heart* 9:113-119.

- [5] Sanchez-Larsen A, García-García J, Ayo-Martin O, Hernández-Fernández F, Díaz-Maroto I, Fernández-Díaz E, Monteagudo M, Segura T (2016) Has the aetiology of ischaemic stroke changed in the past decades? Analysis and comparison of data from current and historical stroke databases. *Neurologia* (16): 30168-2.
- [6] World Health Organization. Projections of mortality and causes of death, 2015 and 2030. Available at: [http://www.who.int/healthinfo/global\\_burden\\_disease/projections/en/](http://www.who.int/healthinfo/global_burden_disease/projections/en/). Date accessed: June 2017.
- [7] Krijthe BP, Kunst A, Benjamin EJ, Lip GY, Franco OH, Hofman A, Witteman JC, Stricker BH, Heeringa J (2013) Projections on the number of individuals with atrial fibrillation in the European Union, from 2000 to 2060. *Eur Heart J* 34(35):2746-2751.
- [8] Marini C, De Santis F, Sacco S, Russo T, Olivieri L, Totaro R, Carolei A (2005) Contribution of atrial fibrillation to incidence and outcome of ischemic stroke: results from a population-based study. *Stroke* 36:1115-1119.
- [9] January CT, Wann LS, Alpert JS, Calkins H, Cigarroa JE, Cleveland JC Jr, Conti JB, Ellinor PT, Ezekowitz MD, Field ME, Murray KT, Sacco RL, Stevenson WG, Tchou PJ, Tracy CM, Yancy CW (2014) American College of Cardiology/American Heart Association Task Force on Practice Guidelines. AHA/ACC/HRS guideline for the management of patients with atrial fibrillation: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and the Heart Rhythm Society. *J Am Coll Cardiol* 64:1-76.
- [10] Camm AJ, Kirchhof P, Lip GYH, Schotten U, Savelieva I, Ernst S, Van Gelder IC, Al-Attar N, Hindricks G, Prendergast B, Heidbuchel H, Alfieri O, Angelini A, Atar D, Colonna P, De Caterina R, De Sutter J, Goette A, Gorenk B, Heldal M, Hohloser SH, Kolh P, Le Heuzey JY, Ponikowski P, Rutten FH (2010) ESC Committee for Practice Guidelines. Guidelines for the management of atrial fibrillation: the Task Force for the Management of Atrial Fibrillation of the European Society of Cardiology (ESC). *Eur Heart J* 31:2369-2429.
- [11] Gladstone DJ, Bui E, Fang J, Laupacis A, Lindsay MP, Tu JV, Silver FL, Kapral MK (2009) Potentially preventable strokes in high-risk patients with atrial fibrillation who are not adequately anticoagulated. *Stroke* 40:235-240.
- [12] Olavarria VV, Delgado I, Hoppe A, Brunser A, Cárcamo D, Díaz-Tapia V, Lavados PM (2011) Validity of the NIHSS in predicting arterial occlusion in cerebral infarction is time-dependent. *Neurology* 76:62-68.
- [13] Maas MB, Furie KL, Lev MH, Ay H, Singhal AB, Greer DM, Harris GJ, Halpern E, Koroshetz WJ, Smith WS (2009) National Institutes of Health Stroke Scale score is poorly predictive of proximal occlusion in acute cerebral ischemia. *Stroke* 40:2988-2993.
- [14] Pisters R, Lane DA, Nieuwlaar R, de Vos CB, Crijns HJ, Lip GY (2010) A novel user-friendly score (HAS-BLED) to assess 1-year risk of major bleeding in patients with atrial fibrillation: The Euro Heart Survey. *Chest* 138(5):1093-1100.
- [15] Suto Y, Kowa H, Nakayasu H, Awaki E, Saito J, Irizawa Y, Nakashima K (2011) Relationship between three-year survival and functional outcome at discharge from acute-care hospitals in each subtype of first-ever ischemic stroke patients. *Intern Med* 50: 1377- 1383.
- [16] Kaasenbrood F, Hollander M, Rutten FH, Gerhards LJ, Hoes AW, Tieleman RG (2016) Yield of screening for atrial fibrillation in primary care with a hand-held, single-lead electrocardiogram device during influenza vaccination. *Europace* 18(10):1514-1520.
- [17] Lowres N, Neubeck L, Redfern J, Freedman SB (2013) Screening to identify unknown atrial fibrillation. A systematic review. *Thromb Haemost* 110(2):213-222.
- [18] Kamiński R, Kosiński A, Brala M, Piwko G, Lewicka E, Dąbrowska-Kugacka A, Raczak G, Kozłowski D, Grzybiak M (2015) Variability of the Left Atrial Appendage in Human Hearts. *PLoS One* 10(11):0141901.
- [19] Chang CJ, Chen YT, Liu CS, Lin WY, Lin CL, Lin MC, Kao CH (2016) Atrial Fibrillation Increases the Risk of Peripheral Arterial Disease With Relative Complications and Mortality: A Population-Based Cohort Study. *Atrial Fibrillation Increases the Risk of Peripheral Arterial Disease With Relative Complications and Mortality. A Population-Based Cohort Study. Medicine (Baltimore)* 95(9):3002.